

EXPORT OF BOVINE SEMEN TO IRAQ

DEPARTMENT FOR ENVIRONMENT, FOOD AND RURAL AFFAIRS SCOTTISH GOVERNMENT

WELSH GOVERNMENT

DEPARTMENT OF AGRICULTURE, ENVIRONMENT AND RURAL AFFAIRS, NORTHERN IRELAND

HEAI	LTH CERTIFICATE	N	o:
EXP	ORTING COUNTRY:	UNITED KINGDOM	
FOR	COMPLETION BY:	OFFICIAL VETERINARIAN	
I.	IDENTIFICATION OF THE	SEMEN	
(na	tity of donor bull(s) mme, date of birth, egistration number)	Breed Number and voluments semen stra	
	* Explanation of date	code(s)	
II.	ORIGIN OF THE SEMEN		
1)	Name and address of Se	emen Collection Centre:	
2)	Approval number of Sen	men Collection Centre:	
3)	Name and address of co	onsignor:	
			• • • • • • • • • • • • • • • • • • • •
4)	The semen consignment	is to be sent from (place o	f loading).
**/		is to be sent from (place o	

- I, the undersigned, certify that:
- the semen described in Paragraph I was derived from bull(s) standing at the Semen Collection Centre described in Paragraph II, which is approved by the competent veterinary authorities of the United Kingdom and is under official veterinary control, being directly supervised by an Authorised Centre Veterinarian;
- 2) prior to entering the approved Semen Collection Centre, all bovine animals have been subjected to a period of isolation of at least 28 days in special accommodation approved by the competent veterinary authority;
- 3) within the 28 day period prior to commencing the period of isolation referred to in paragraph IV 2) above, all the bovine animals have been subjected to the following tests:
- b) TUBERCULOSIS: a comparative intradermal tuberculin test using avian and bovine purified protein derivative (PPD) tuberculins, with negative results according to the Department's standard interpretation;
- c) BOVINE VIRAL DIARRHOEA/MUCOSAL DISEASE (BVD-MD): (i) a virus isolation test or a test for virus antigen, with negative results. In the case of an animal less than six months old, the test has been deferred until that age has been reached;
 - (ii) a serological test to determine the serological status of every animal;
- d) INFECTIOUS BOVINE RHINOTRACHEITIS/INFECTIOUS PUSTULAR VULVO-VAGINITIS
 (IBR/IPV): a * serum neutralisation test (SNT) or * enzyme-linked
 immunosorbent assay (ELISA) with negative results;
- e) ENZOOTIC BOVINE LEUKOSIS: * an agar gel immunodiffusion test (AGIDT) or * enzyme-linked immunosorbent assay (ELISA), with negative results;
- 4) during the period of 28 days isolation referred to in paragraph IV 2) above, all the bovine animals have been subjected to the following tests:

- b) BOVINE VIRAL DIARRHOEA/MUCOSAL DISEASE (BVD-MD):(i) a virus isolation test or a test for virus antigen, with negative results;
 - (in) a serological test to determine the serological status of every animal;
- BOVINE GENITAL CAMPYLOBACTERIOSIS (Campylobacter fetus venerealis): *an immunofluorescent antibody test * or a culture test on a sample of preputial material, with negative results;
- d) TRICHOMONIASIS (Trichomonas fetus): a microscopic examination and culture test on a sample of preputial material, with negative results;
- e) INFECTIOUS BOVINE RHINOTRACHEITIS/INFECTIOUS PUSTULAR VULVO-VAGINITIS (IBR/IPV): * a serum neutralisation test (SNT) * or enzyme-linked immunosorbent assay (ELISA) with negative results;
- 5) at regular 12-monthly intervals, all the bovine animals at the approved Semen Collection Centre are subjected to the following tests:
- b) TUBERCULOSIS: a comparative intradermal tuberculin test using avian and bovine purified protein derivative (PPD) tuberculins, with negative results according to the Department's standard interpretation;
- c) BOVINE VIRAL DIARRHOEA/MUCOSAL DISEASE (BVD-MD): a serological test on all animals negative to previous serological tests, with negative results. If any animal has become serologically positive, every ejaculate of that animal collected since the last negative test has been either discarded or tested for virus with negative results;
- d) BOVINE GENITAL CAMPYLOBACTERIOSIS (Campylobacter fetus venerealis): * an immunofluorescent antibody test * or a culture test on a sample of preputial material, with negative results;
- e) TRICHOMONIASIS (*Trichomonas fetus*): a microscopic examination and culture test on a sample of preputial material, with negative results;
- f) INFECTIOUS BOVINE RHINOTRACHEITIS/INFECTIOUS PUSTULAR VULVO-VAGINITIS (IBR/IPV): * a serum neutralisation test (SNT) * or enzyme-linked immunosorbent assay (ELISA) with negative results;
- g) ENZOOTIC BOVINE LEUKOSIS: * an agar gel immunodiffusion test (AGIDT) * or enzyme linked-immunosorbent assay (ELISA), with negative results;
- 6) regarding BLUE TONGUE VIRUS (BTV), the semen was obtained from donor bull(s) which comply with at least one of the following conditions:
 - *(a) they were kept in a BTV free country or zone for a period of at least 60 days before commencement of, and during, collection of the semen; OR
 - *(b) they were subjected to a serological test according to the OIE Terrestrial Manual to detect antibodies to the BTV group, with negative results, at least every 60 days during the collection period and between 28 and 60 days after the final collection for this consignment; OR
 - \star (c) they were subjected, with negative results, to an agent identification test for BTV according to the OIE Terrestrial Manual carried out on blood

samples collected:

- (i) at commencement and final collection of the semen for this consignment, and
- (ii) during the period of semen collection for this consignment:
 - *i. at least every seven days, in the case of a virus isolation test, or *ii. at least every 28 days, in the case of a polymerase chain reaction (PCR) test;
- regarding BOVINE VIRAL DIARRHOEA/MUCOSAL DISEASE (BVD-MD): prior to the initial despatch of semen from BVD-MD serologically positive bulls, a semen sample from each animal has been subjected to a virus isolation or virus antigen test for BVD. In the event of a positive test, the bull was removed from the centre and all of its semen destroyed;
- all the above tests have been carried out at laboratories approved by the competent veterinary authority;
- 9) the semen described above is derived from donor bulls which:
 - a) showed no clinical signs of disease on the day(s) of semen collection;
 - b) have not been used for natural service while at the approved Semen Collection Centre;
 - c) have not been vaccinated against foot and mouth disease;
- 10) the semen described above:
 - a) has been treated during processing by the addition of antibiotics to produce these concentrations in the final diluted semen:
 - *EITHER: not less than 500 µg per ml streptomycin, 500 IU per ml penicillin, 150 µg per ml lincomycin, 300 µg per ml spectinomycin

*OR:																							an	е	qu:	iva	ale	ent	
effe	ct	aga	ins	t (Cam	pyl	oba	ct	ers	3,	Lep	oto	sp	lre	s	and	Му	700	p1	asn	nas	,							
name	ly.											• • •	••,				٠,٠.	/ .						٠.					
												. 🔍			• •	,	/	•											

- b) has been stored in an approved Semen Collection Centre for at least 30 days before dispatch;
- c) has been placed in a previously cleaned and disinfected container, which before dispatch was sealed under veterinary supervision. The seal number is stated at paragraph III 3) above;
- 11) the United Kingdom is free from foot and mouth disease and rinderpest in accordance with the OIE Terrestrial Animal Health Code;
- 12) the approved Semen Collection Centre in which the semen described above in paragraph I was collected is located at the centre of an area of 10 kilometres radius in which there has been no case of foot and mouth disease for at least 30 days prior to the date(s) of semen collection.

*	Delete	as	appropriate
---	--------	----	-------------

Official Stamp	Signed RCVS Official Veterinarian
	Name in block letters
Date	Address